

Original Article

Surgical outcomes of patients with corrosive ingestion: a retrospective analysis

Farhan Majeed ^a, Ahmed Raza ^{b,*}, Muhib Ullah ^a, Farhan Ullah ^a, Osama Zafar ^c, Nabeela Farhan ^d

^a Department of Thoracic Surgery, Combined Military Hospital Rawalpindi, Pakistan

^b Department of Thoracic Surgery, Combined Military Hospital Peshawar, Pakistan

^c Queen Elizabeth Hospital Birmingham, United Kingdom

^d Al Shifa Home Health, Rawalpindi, Pakistan

* Correspondence: ahmedraza1470@gmail.com



Citation: Majeed F, Raza A, Ullah M, Ullah F, Zafar O, Farhan N. Surgical outcomes of patients with corrosive ingestion: a retrospective analysis. J Basic Clin Med Sci. 2024;3:27-33.

Received: 12 October 2024

Revised: 18 December 2024

Accepted: 20 December 2024

Published: 23 December 2024

Publisher's Note: Logixs Journals remains neutral concerning jurisdictional claims in its published subject matter, including maps and institutional affiliations.



Copyright: © 2024 The Author(s).

This is an open access article distributed under the terms of the [Creative Commons Attribution \(CC BY\) License](https://creativecommons.org/licenses/by/4.0/). The use, distribution, or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Abstract

Corrosive ingestion can cause severe complications, including perforation, stricture, and fistula, which often require surgical intervention to prevent further damage. This retrospective observational study analyzed surgical outcomes and gender differences in ingestion types (accidental vs. suicidal) and causative substances (acid, alkali, ammonia, detergent, or mixed) among 99 patients at a tertiary care hospital in Rawalpindi from January 2002 to January 2024. The average age of the patients was 30.36 ± 10.18 years, with 61.62% female patients and 38.38% male patients. Accidental ingestion was more common in males (94.74%), whereas suicidal ingestion was more common in females (21.31%, $p = 0.026$). Mixed substances were the most common substances ingested, particularly among females (60.66%, $p = 0.048$). All patients underwent upper gastrointestinal endoscopy, and diagnostic laparoscopy or feeding jejunostomy was the most common surgical procedure (57.58%). The esophagus (68.69%) was the most affected site. Surgical site infections (16.16%) and strictures (11.11%) were the most frequent complications, with a mortality rate of 4.04%. Our study concluded that timely surgical management could reduce the risk of complications and enhance patient outcomes. Corrosive ingestion often affects multiple areas of the gastrointestinal tract that require several surgical interventions. In certain cases, complex procedures, such as colon interposition and total laryngopharyngoesophagectomy (TLPO), which require skilled surgeons as well as advanced management centers, are performed to manage corrosive injuries. Therefore, improving these patients' outcomes via interventions focused on training in complex surgical procedures is crucial. However, more research is needed to identify the most effective approaches for the surgical management of corrosive ingestion.

Keywords

Corrosive ingestion; Surgical management; Complications; Retrospective analysis

1. Introduction

Corrosive ingestion refers to the intentional or accidental ingestion of corrosive substances, such as acids or alkalis, which can cause severe damage to the digestive tract. This can lead to serious complications, including perforation of the esophagus and stomach. Corrosive ingestion is a major public health issue globally [1,2]. It is more prevalent in developing countries, but it is also common in developed countries. Even in the United States, approximately 5,000–15,000 corrosive ingestions are reported annually [3]. Annually, more than 40,000 cases of corrosive ingestion are reported in England and Wales.

In developing countries such as Pakistan, it is very prevalent, although most cases are not reported due to cultural barriers [4]. The lack of safe "childproof" containers, un-

regulated access to corrosive agents, a culturally specific inclination to ingest corrosives for suicidal purposes, and the absence of standard medical care in rural areas [4,5,6]. The medical literature concerning the epidemiology and clinical presentation of corrosive ingesting patients is inadequate. The World Health Organization (WHO) reported that in the past 17 years, only 37 papers concerning corrosive ingestion in low- and lower-middle-income countries have been published [7].

Surgical intervention plays a crucial role in the management of corrosive ingestion patients, particularly in cases where severe damage to the gastrointestinal tract has occurred [8,9]. The type of surgical intervention depends on the damaged area, and multiple surgical approaches can be used in multiple affected areas [9]. Surgical evaluation is essential for assessing the extent of damage caused by corrosive substances [10]. This may involve various diagnostic procedures, such as endoscopy, imaging studies (e.g., CT scans), and clinical assessments, to determine the severity and location of the injury. Corrosive ingestion can lead to a range of complications, including perforation, stricture, and fistula in the gastrointestinal tract [11]. Surgical intervention may be necessary to address these complications, such as repairing perforations, dilating strictures, or performing bowel resections in severe cases. Surgical intervention aims to prevent further damage to the gastrointestinal tract and surrounding structures. Therefore, the present study was conducted to retrospectively analyze the surgical outcomes of patients managed surgically for corrosive ingestion. Moreover, sex differences in ingestion types (accidental and suicidal) and ingested causative substances (acid, alkali, ammonia, detergents, and mixed) were assessed in corrosive ingestion cases.

2. Methods

2.1. Study design

This was a retrospective observational study that assessed the surgical outcomes of patients who ingested corrosive substances.

2.2. Study duration

Data from January 2002 to January 2024 were retrieved from the database and patients' medical records at a tertiary care hospital in Rawalpindi.

2.3. Ethical considerations

The study adhered to ethical guidelines and received all necessary approvals. The study obtained ethical approval from the Ethical Committee/Institutional Review Board, Combined Military Hospital, Rawalpindi (No. 267).

2.4. Sample size

The sample size was calculated using the proportion of patients with corrosive ingestion injuries who underwent surgical procedures (6.2%) at a 95% confidence level and a 5% margin of error [12,13]. The final sample size was calculated as 90, which was further increased to 99 to strengthen the study findings.

2.5. Selection criteria

The study included admitted patients who underwent surgical intervention for corrosive ingestion and who were 5 years of age or older, with documented corrosive ingestion durations of 3 months or longer, and who were receiving surgical management. Patients who presented with malignant or other benign strictures were excluded. In addition, all patients whose medical records were incomplete were excluded from the study.

2.6. Statistical analysis

The data were analyzed via SPSS (version 27.0) by calculating frequencies, percentages, means, and standard deviations (SDs). The chi-square test was applied to assess the gender-wise comparison of ingestion types and causative substances in corrosive ingestion cases. The results were considered significant at $p < 0.05$.

3. Results

Table 1 shows that the average age of the patients was 30.36 ± 10.18 years, with 61.62% female patients and 38.38% male patients. Most of the cases (84.85%) were accidental, whereas 15.15% were suicidal ingestion cases. Among the 99 patients, 72.73% had no comorbidities. Among the ingested substances, a mixed combination (55.56%) was most common, followed by acid ingestion (25.25%).

Table 1. Sociodemographic indicators and medical history of corrosive ingestion cases (n = 99).

Variables		Frequency (%)	Mean \pm Std. Deviation
Age (in years)		-	30.36 \pm 10.18
Gender	Male	38 (38.38)	-
	Female	61 (61.62)	-
Comorbidity	Hypertension	15 (15.15)	-
	Diabetes	2 (2.02)	-
	Diabetes, hypertension	7 (7.07)	-
	Any other	3 (3.03)	-
	No comorbidity	72 (72.73)	-
Ingestion	Accidental	84 (84.85)	-
	Suicidal	15 (15.15)	-
Ingested causative substance	Acid	25 (25.25)	-
	Detergents	9 (9.09)	-
	Alkali	5 (5.05)	-
	Ammonia	5 (5.05)	-
	Mixed	55 (55.56)	-

Table 2 shows that accidental ingestion was more common among males (94.74%) than females (78.69%), whereas suicidal ingestion was significantly greater among females (21.31%) than males (5.26%) ($p = 0.026$). Moreover, compared with males, most of the females (60.66%) ingested mixed substances, while 29.51% ingested acid (47.37% and 18.42%, respectively), and these differences were statistically significant ($p = 0.048$).

Table 2. Sexwise comparison of ingestion types and causative substances in corrosive ingestion cases (n = 99).

Variables		Gender		Degree of Freedom (df)	Chi-Square Value (X ²)	p value *
		Male Frequency (%)	Female Frequency (%)			
Ingestion	Accidental	36 (94.74)	48 (78.69)	1	4.691	0.026 *
	Suicidal	2 (5.26)	13 (21.31)			
Ingested causative substance	Acid	7 (18.42)	18 (29.51)	4	9.577	0.048 *
	Alkali	3 (7.89)	2 (3.28)			
	Ammonia	4 (10.53)	1 (1.64)			
	Detergents	6 (15.79)	3 (4.92)			
	Mixed	18 (47.37)	37 (60.66)			

* Variables were compared using the chi-square test. ** Significant value ($p \leq 0.05$).

Table 3 shows that all patients underwent upper gastrointestinal endoscopy, whereas contrast studies were performed in 97.98% of patients. Moreover, diagnostic laparoscopy or feeding jejunostomy was the most common surgical procedure (57.58%), followed by complete esophagectomy with gastric pull-up (34.34%). Furthermore, complex surgeries, such as esophagogastrectomy with colonic interposition and Roux-en-Y, were performed in 21.21% of patients. The most affected site of injury was the esophagus (68.69%), followed by the pylorus (36.36%) and gastroesophageal junction (23.23%). Among the complications, surgical site infections or abscesses were the most common (16.16%), followed by stricture formation (11.11%) and respiratory complications such as empyema, pneumonia, or atelectasis (10.10%). Mortality was recorded in 4.04% of the 99 patients.

Table 3. Procedures, injuries, management, and complications in corrosive ingestion cases.

Variables	Frequency (%)
Evaluation procedures	Upper gastrointestinal endoscopy
	99 (100.00)
	Contrast study
Injury location/extent	18 (18.18)
	Esophagus
	68 (68.69)
	Pylorus
	36 (36.36)
	Gastroesophageal junction
	23 (23.23)
	Proximal stomach
Surgical procedures	19 (19.19)
	Distal stomach
	17 (17.17)
	Pharynx/hypopharynx
	2 (2.02)
	Diagnostic laparoscopy/feeding jejunostomy
	57 (57.58)
	Complete esophagectomy with gastric pull-up
	34 (34.34)
	Pyloroplasty
Complications	25 (25.25)
	Esophago-gastrectomy, colonic interposition, Roux-en-Y
	21 (21.21)
	Gastrojejunostomy
	7 (7.07)
	Partial esophagectomy with gastric pull-up
	6 (6.06)
	Tracheoesophageal fistula repair
	3 (3.03)
	Total laryngopharyngoesophagectomy
	2 (2.02)
	Esophagectomy, colonic interposition, Roux-en-Y
	1 (1.01)
	Abscess/surgical site infections
	16 (16.16)
	Stricture formation
	11 (11.11)
	Empyema/pneumonia/atelectasis
	10 (10.10)
	Cardiac arrhythmias
	8 (8.08)
	Recurrent laryngeal nerve injury
	7 (7.07)
	Hemorrhage
	5 (5.05)
	Anastomotic leak
	4 (4.04)
	Mortality
	4 (4.04)
	Conduit necrosis
	3 (3.03)

4. Discussion

The current study reported the surgical management of patients who ingested corrosive substances. Most of the cases were accidental, and mixed substances were reported as the most commonly ingested materials. The majority of the patients did not have comorbidities. Moreover, sex differences were significant in terms of ingestion patterns as well as the types of substances ingested. All patients underwent upper gastrointestinal endoscopy, and diagnostic laparoscopy or feeding jejunostomy was the most frequently performed surgical procedure. The esophagus was the most affected site of in-

jury, followed by the pylorus and gastroesophageal junction. Furthermore, postoperative complications were common, with surgical site infections and strictures being the most commonly reported, whereas mortality and anastomotic leaks were rare.

Corrosive ingestion triggers full-thickness esophagogastric necrosis, which represents the most severe form of injury to the alimentary tract. The degree of severe damage is typically associated with the ingestion of large quantities of concentrated corrosive agents, often in cases of suicidal intent, as observed in most cases in our study. The injuries are generally confined to the esophagus and stomach, with the duodenum being relatively spared. This sparing is likely owing to pyloric spasm triggered by the corrosive substance or the alkaline environment of the duodenum. Severe metabolic complications, including acidosis and dehydration, often accompany such injuries. Furthermore, disruption of the gut mucosal barrier, in addition to contamination of the peritoneal and mediastinal spaces, often leads to life-threatening sepsis.

The management of corrosive ingestion requires immediate resuscitation to correct fluid, electrolyte, and acid-base imbalances, along with the administration of broad-spectrum antibiotics and timely surgical intervention. Significant differences in mortality rates were observed between the surgical group (67%) and the supportive treatment group (100%) in a comparative study involving 27 patients [14]. In contrast, our study reported a mortality rate of 4%, which may be attributed to effective management strategies, the extent of injury, and the retrospective design of our study, which did not include long-term patient follow-up. In such cases, surgical treatment generally focuses on excising nonviable tissue and ensuring effective peritoneal and mediastinal drainage. There is a dearth of literature on the management and outcomes of severe esophagogastric injuries owing to corrosive ingestion. Cattani et al., in their review, reported immediate postoperative mortality in 2 out of 9 patients, with further deaths occurring in the months after surgical resection [15]. Moreover, a review on the outcome of gastric perforation resulting from corrosive ingestion revealed that patients with gastric necrosis and perforation had poor survival outcomes, whereas those with isolated perforation that was repaired primarily experienced improved survival [16].

Higher mortality rates have been reported when a thoracoabdominal approach is employed for esophageal resection following severe corrosive injuries. In a study by Lai et al., 18 patients who underwent emergency surgery were recruited, and the overall mortality rate was reported to be 66.7%, with four deaths occurring among patients who underwent esophagectomy and stomach resection via the thoracoabdominal method [14]. In contrast, mortality was lower (37.5%) in patients who underwent esophageal stripping with stomach resection through the abdominal route. These findings suggest that esophageal stripping is a safer alternative to the thoracoabdominal approach in such cases.

Despite the severity of injuries and the extensive surgical resection often needed, the perioperative mortality rate reported in the literature is relatively low [17]. Factors associated with improved outcomes include aggressive resuscitation, timely surgical intervention to limit further injury, complete excision of necrotic tissue, transhiatal esophageal resection, and postoperative nutritional support provided through jejunostomy feeding [18]. Additionally, corrosive-induced periesophageal edema has been suggested to aid in transhiatal resection by reducing procedural complexity and minimizing morbidity [19].

Our study findings revealed significant gender-based differences in patterns of substance ingestion. Accidental ingestion was more prevalent among males, which may be linked to occupational or environmental exposures associated with male-dominated activities [20,21,22]. Conversely, suicidal ingestion occurs more frequently among females,

which may be due to psychological and sociocultural factors, as females may experience higher levels of emotional distress and societal pressures that contribute to self-harm [23,24]. Additionally, females were more likely to ingest mixed substances, whereas males were more likely to ingest acid. These findings highlight the need for tailored prevention strategies, such as promoting workplace safety for males and providing targeted mental health support for females, to address the specific needs and underlying factors associated with these behaviors [25,26]. To explore related qualitative factors and determine the contributing sociodemographic parameters unaddressed by the current study, further research is needed, which is a potential weakness of this study.

5. Conclusions

Our study concluded that timely surgical management could reduce the risk of complications and enhance patient outcomes. Corrosive ingestion often affects multiple areas that require several surgical interventions. In certain cases, complex procedures, such as colon interposition and total laryngopharyngoesophagectomy (TLPO), which require skilled surgeons as well as advanced management centers, are performed to manage corrosive injuries. Therefore, improving these patients' outcomes via interventions focused on training in complex surgical procedures is crucial. Further research is needed to identify the most effective approaches for the surgical management of corrosive ingestion.

Author contributions: Conceptualization, FM, AR, MU, FU, OZ and NF; methodology, FM, AR, MU, FU, and NF; software, AR, OZ, and NF; validation, FM, MU, FU and OZ; formal analysis, AR, OZ, and NF; investigation, AR, MU, FU, and OZ; resources, FM, and AR; data curation, AR, OZ, and NF; writing—original draft preparation, MU, FU, OZ and NF; writing—review and editing, FM, and AR; visualization, FM, MU, FU and OZ; supervision, FM, and AR; project administration, AR, MU, FU, and NF. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no specific grant from the public, commercial, or not-for-profit funding agencies.

Ethics statement: The study was approved by the Ethical Committee/Institutional Review Board, Combined Military Hospital, Rawalpindi (No. 267).

Consent to participate: Not applicable.

Data availability: The data supporting this study's findings are available from the corresponding author, Ahmed Raza, upon reasonable request.

Acknowledgments: None.

Conflicts of interest: The authors declare no conflicts of interest.

References

- [1] Kalayarasan R, Ananthakrishnan N, Kate V. Corrosive ingestion. *Indian J Crit Care Med*. 2019;23(Suppl 4):S282-6. <https://doi.org/10.5005/jp-journals-10071-23305>
- [2] Challine A, Maggiori L, Katsahian S, Corté H, Goere D, Lazzati A, et al. Outcomes associated with caustic ingestion among adults in a national prospective database in France. *JAMA Surg*. 2022;157(2):112–9. <https://doi.org/10.1001/jamasurg.2021.6368>
- [3] Mahmoudvand, Z., Shanbehzadeh, M., Shafiee, M. Kazemi-Arpanahi H. Developing the minimum data set of the corrosive ingestion registry system in Iran. *BMC Health Serv Res*. 2022;22:1207. <https://doi.org/10.1186/s12913-022-08576-0>
- [4] Ain QU, Jamil M, Safian HA, Akhter TS, Batool S, Arshad M, et al. Assessing the degree of acute esophageal injury secondary to corrosive intake: insights from a public sector hospitals of a developing country. *Cureus*. 2020;12(10):e10858. <https://doi.org/10.7759/cureus.10858>
- [5] Özcan C, Ergün O, Şen T, Mutaf O. Gastric outlet obstruction secondary to acid ingestion in children. *J Pediatr Surg*. 2004;39(11):1651-3. <https://doi.org/10.1016/j.jpedsurg.2004.07.008>
- [6] Grabska K, Pilarska I. Acute poisoning among children and adolescents: a narrative review. *Med Sci Pulse*. 2022;16(3):33-9. <https://doi.org/10.5604/01.3001.0015.9656>

- [7] Abbasi HW, Aftab W, Shahid S, Ali M, Zahid J, Naz S, et al. Epidemiological characteristics, spectrum of upper gastrointestinal tract injuries and burden of corrosive intake patients in a tertiary care hospital in Islamabad. *Life Sci.* 2023;4(2):6-9. <https://doi.org/10.37185/lnS.1.1.287>
- [8] Gupta V, Shah J, Yadav TD, Kumar P, Wig JD, Kochhar R. Emergency surgical intervention in acute corrosive ingestion: single-center experience from India. *ANZ J Surg.* 2023;93(12):2864-9. <https://doi.org/10.1111/ans.18576>
- [9] Agarwal A, Srivastava DN, Madhusudhan KS. Corrosive injury of the upper gastrointestinal tract: the evolving role of a radiologist. *Br J Radiol.* 2020;93(1114):20200528. <https://doi.org/10.1259/bjr.20200528>
- [10] Yu CH, Su YJ, Lai YC. Fatal Zargar grade 3b corrosive injury after hydrochloric acid ingestion: a case report. *Medicine.* 2024;103(40):e40017. <https://doi.org/10.1097/MD.00000000000040017>
- [11] Tettey M, Edwin F, Tamatey M, Okonta K, Entsua-Mensah K, Adzamlı I, et al. Management of caustic pharyngoesophageal injuries in Korle Bu Teaching Hospital, Accra, Ghana: a 12-year retrospective review of 29 cases. *Pan Afr Med J.* 2022;42:213. <https://doi.org/10.11604/pamj.2022.42.213.30206>
- [12] Javed A, Pal S, Krishnan EK, Sahni P, Chattopadhyay TK. Surgical management and outcomes of severe gastrointestinal injuries due to corrosive ingestion. *World J Gastrointest Surg.* 2012;4(5):121-5. <http://dx.doi.org/10.4240/wjgs.v4.i5.121>
- [13] Open EPI. Sample size. 2024 [cited 17 December 2024]. Available from: <https://www.openepi.com/SampleSize/SSPropor.htm>.
- [14] Lai KH, Huang BS, Huang MH, Huang MS, Wu JK, Liu M, et al. Emergency surgical intervention for severe corrosive injuries of the upper digestive tract. *Zhonghua Yi Xue Za Zhi (Taipei).* 1995;56(1):40-6.
- [15] Cattani P, Munoz-Bongrand N, Berney T, Halimi B, Sarfati E, Celerier M. Extensive abdominal surgery after caustic ingestion. *Ann Surg.* 2000;231(4):519-23. <https://doi.org/10.1097/00000658-200004000-00010>
- [16] Ceylan H, Ozokutan BH, Gündüz F, Gözen A. Gastric perforation after corrosive ingestion. *Pediatr Surg Int.* 2011;27:649-53. <https://doi.org/10.1007/s00383-010-2739-6>
- [17] Scriba MF, Kotze U, Naidoo N, Jonas E, Chinnery GE. Aorto-oesophageal fistula after corrosive ingestion: a case report. *World J Gastrointest Surg.* 2022;14(5):506-13. <https://doi.org/10.4240/wjgs.v14.i5.506>
- [18] Ríos G, Rodríguez L, Lucero Y, Miquel I, Arancibia ME, Alliende F. Endoscopic findings associated with button battery ingestion in children: do we need to change the protocol for managing gastric location? *Pediatr Emerg Care.* 2020;36(11):523-6. <https://doi.org/10.1097/pec.0000000000001415>
- [19] Patterson KN, Beyene TJ, Gil LA, Minneci PC, Deans KJ, Halaweish I. Procedural and surgical interventions for esophageal stricture secondary to caustic ingestion in children. *J Pediatr Surg.* 2023;58(9):1631-9. <https://doi.org/10.1016/j.jpedsurg.2023.01.048>
- [20] Awogbami SO, Ogunyemi O, Adebayo PA, Raimi MO. Protecting the health of black communities: assessing the impact of environmental hazards from gold mining activities on health outcomes among residents of Osun State, Nigeria. *J Med Internet Res.* Forthcoming 2024. Available from: <https://preprints.jmir.org/preprint/66508>
- [21] Kwesi SS, Dartey E, Kuffour RA, Dekugmen DY, Osei-Bonsu R, Mintah DK, et al. Occupational health and safety practices among small-scale mining workers in Ghana. *Eur J Sci Innov Technol.* 2024;4(2):26-39.
- [22] Orellana C, Kreshpaj B, Burstrom B, Davis L, Frumento P, Hemmingsson T, et al. Organisational factors and under-reporting of occupational injuries in Sweden: a population-based study using capture-recapture methodology. *Occup Environ Med.* 2021;78(10):745-52. <https://doi.org/10.1136/oemed-2020-107257>
- [23] Aktar S, Tribe R. The experience of self-harming behaviours that inflict external injuries to the body in UK-based Bangladeshi, Indian and Pakistani women: a literature review. *Int Rev Psychiatry.* 2024(4-5):442-50. <https://doi.org/10.1080/09540261.2024.2306620>
- [24] Abbas N, Ali WB, Qandeel, Gull S, Begum S. Exploring socio-cultural determinants of female suicides: a qualitative study of Gahkuch, Gilgit-Baltistan. *Soc Sci Rev Arch.* 2025;3(1):450-60. <https://doi.org/10.70670/sra.v3i1.323>
- [25] Arensman E, O'Connor C, Leduc C, Griffin E, Cully G, Ní Dhálaigh D, et al. Mental health promotion and intervention in occupational settings: protocol for a pilot study of the MENTUPP intervention. *Int J Environ Res Public Health.* 2022;19(2):947. <https://doi.org/10.3390/ijerph19020947>
- [26] Pirkis J, Bantjes J, Dandona R, Knipe D, Pitman A, Robinson J, et al. Addressing key risk factors for suicide at a societal level. *Lancet Public Health.* 2024;9(10):e816-24. [https://doi.org/10.1016/S2468-2667\(24\)00158-0](https://doi.org/10.1016/S2468-2667(24)00158-0)